

Mini-Mental State Examination의 한국 노인 정상기준 연구*

이동영¹⁾ · 이강욱²⁾ · 이정희²⁾ · 김기웅¹⁾ · 주진형³⁾
 윤중철⁴⁾ · 김성운⁵⁾ · 우성일⁶⁾ · 우종인⁴⁾†

A Normative Study of the Mini-Mental State Examination
 in the Korean Elderly*

Dong Young Lee, M.D., Ph.D.,¹⁾ Kang Uk Lee, M.D., Ph.D.,²⁾ Jung Hie Lee, M.D., Ph.D.,²⁾
 Ki Woong Kim, M.D., Ph.D.,¹⁾ Jin Hyeong Jhoo, M.D.,³⁾ Jong Chul Youn, M.D.,⁴⁾
 Sung Youn Kim, M.D., Ph.D.,⁵⁾ Sung Il Woo, M.D., Ph.D.,⁶⁾ Jong Inn Woo, M.D., Ph.D.⁴⁾†

국문초록

연구목적 :

MMSE in the Korean version of the CERAD assessment packet(MMSE - KC)
 MMSE - KC

방법 :

60~90 618 MMSE - KC MMSE - KC
 CERAD - K 가 가 ,
 가
 MMSE - KC

: 2002 1 7

: 2002 4 8

1996 1997

(Grant No : HMP - 96 - M - 2 - 0029)

2000

(Grant No : 04 - 2000 - 045)

1)

Department of Neuropsychiatry, Kyunggi Provincial Hospital for the Elderly, Yongin

2)

Department of Neuropsychiatry, Kangwon National University College of Medicine, Chunchon

3)

Department of Neuropsychiatry, Anyang Metro Hospital, Anyang

4)

Department of Psychiatry, Seoul National University College of Medicine & Clinical Research Institute, Seoul National University Hospital, Seoul

5)

Department of Psychiatry, Asan Medical Center, Seoul

6)

Department of Psychiatry, Soon Chun Hyang University Hospital, Seoul

†Corresponding author

MMSE - KC

결 과 :

1) 60 MMSE - KC 가 가 .

가 MMSE - KC , 60 ,

2) MMSE - KC , 24

MMSE - KC 5, 25, 75 ,

3) MMSE - KC ,

가

결 론 :

MMSE - KC

가 MMSE - KC .

중심 단어 : MMSE - KC · MMSE

서 론

14)

10)

15)16) MMSE

1975 Folstein , , , , ,

(Mini - Mental State Examination ; MMSE) MMSE 가 ,

가 17)18) 16)19 - 21) 16)

1 - 3) MM- 22)23)

SE

가 ,

MMSE

4) 5) 6)

가

7)8)

MMSE , , , , ,

(percentile) (norm)

9 - 13) MMSE . MMSE

가

MMSE 24)

5)

2. 검사도구 : MMSE-KC

Lee CERAD - K CE-
 RAD 가 MMSE 10 MMSE - KC 가
 , MMSE - KC(MMSE in the Ko-
 rean version of the CERAD assessment packet)
 30) CE-
 RAD 가 MMSE(CERAD - MMSE)
 Folstein (1975) MMSE

가
 Folstein MMSE
 MMSE - KC CERAD - MMSE
 ,
 MMSE - K⁵⁾
 ,
 MMSE - K
 가
 (),
 (, ,) MMSE - K

MMSE -
 K
 , MMSE - KC Folstein MMSE CE-
 RAD - MMSE
 , Folstein MMSE
 subtraction) 가 (serial 2
 (WORLD) , CERAD - MMSE
 , MMSE -
 KC 3가
 , CERAD - MM-
 SE . MMSE - KC

3. MMSE-KC의 시행

MMSE - KC CERAD - K 가 , 1 2 가
 (R²)
)
 MMSE - KC 가
 , Lee CERAD - K . 60

4. MMSE-KC 검사 점수에 영향을 미치는 변인 분석

가
 ,
 MMSE - KC
 (univariate analysis)

(multivariate analysis)
 Pearson
 t - test
 (multiple linear regression)
 1(stage 1)

3가
 , 2(stage 2)
 0.05
 가

가 , 1 2 가
 (R²)
 . 60 가
 (60)
 . 60

50 60 ()
 16)
 가 30
 140/90mmHg
 33) , 가
 1
 , 6
 가
 5. 정상 기준 값의 산출 및 산출된 기준의 타당도 검증
 1) 정상기준 산출
 5 (percentile), 25
 (lower quartile), , 75 (upper qu-
 artile), ,
 2) 기준 타당도 검증

Table 1. Demographic and clinical characteristics of the subjects participating in the normative study of the MMSE-KC

	Men	Women	Total
Number	209	409	618
Age(yr)	73.6 ± 6.3 [†]	71.2 ± 6.0	72.0 ± 6.3
60 - 64	6(2.9) [‡]	48(11.7)	54(8.7)
65 - 69	61(29.2)	143(35.0)	204(33.0)
70 - 74	58(27.8)	103(25.2)	161(26.1)
75 - 79	45(21.5)	71(17.4)	116(18.8)
80 - 84	27(12.9)	29(7.1)	56(9.1)
85 - 89	10(4.8)	14(3.4)	24(3.9)
90	2(1.0)	1(0.2)	3(0.5)
Education(yr)	7.8 ± 5.4	4.8 ± 4.0	5.9 ± 4.8
0	37(17.7)	125(30.6)	162(26.2)
1 - 3	8(3.8)	39(9.5)	47(7.6)
4 - 6	68(32.5)	159(38.9)	227(36.7)
7 - 9	26(12.4)	40(9.8)	66(10.7)
10 - 12	29(13.9)	35(8.6)	64(10.4)
13	41(19.6)	11(2.7)	52(8.4)
Previous rural residence	79(37.8)	224(54.8)	303(49.0)
Hypertension	90(43.1)	162(39.6)	252(40.8)
Diabetes	17(8.1)	49(12.0)	66(10.7)
Alcohol users (lifetime)	129(61.7)	27(6.6)	156(25.2)
Alcohol user (current)	92(44.0)	24(5.9)	116(18.8)
Smoker (lifetime)	133(63.6)	34(8.3)	167(27.0)
Smokers (current)	81(38.8)	24(5.9)	105(17.0)

† : Mean±S.D ‡ : Number(percent)

() 60
 61 가
 가가 MMSE - KC
 (DSM -)
 가 5
 5
 kappa . McKhann
 5
 61 34)
 .

6. 통계 분석 프로그램
 SPSS version
 9.0 for windows

연구 결과

1. 연구대상자의 특성
 1 . 618
 72.0 ± 6.3 , 73.6 ±
 6.3 , 71.2 ± 6.0
 (t=4.6, p<0.01).
 65~79 가

(78.5%; 77.6%),
 ($\chi^2=22.8$, $df=6$, $p<0.01$).
 5.9 ± 4.8 ,
 7.8 ± 5.4 4.8 ± 4.0
 (t=7.8, $p<0.01$).
 30.6% 17.7%
 ($\chi^2=$
 68.0, $df=5$, $p<0.01$).
 가
 49.0% , 37.8%, 54.8%
 가
 ($\chi^2=15.3$, $df=1$, $p<0.01$).
 40.8%, 10.7%
 ,
 61.7%, 6.6% ($\chi^2=219.8$, $df=1$, $p<0.01$),
 44.0%, 5.9% ($\chi^2=129.6$, $df=$
 1, $p<0.01$) 가
 63.6%,
 8.3% ($\chi^2=211.9$, $df=1$, $p<0.01$),
 38.8%, 5.9% ($\chi^2=103.8$, $df=1$, $p<0.01$)
 가
 30 (4.9%),
 가 588 (95.1%)

2. MMSE-KC 점수에 영향을 미치는 변인 분석결과

1) 단변인분석 결과

MMSE - KC
 (r= -0.41, $p<0.01$)
 (r=0.55, $p<0.01$) 가 , MM-
 SE - KC 가
 (t=6.49, $p<0.01$), (t=2.99, $p<0.01$)
 (t=2.76, $p<0.01$) 가 가
 가 MMSE - KC

MMSE - KC

2) 다중 회귀 분석 결과

1(stage 1) , , 3가

Table 2. Multiple linear regression of age, education, sex and other variables on the MMSE-KC score

	SE()	R2	p
Stage 1		0.41*	
Age	-0.21 0.02		0.00064*
Education	0.34 0.03		0.00063*
Sex	1.32 0.28		0.00030*
Stage 2		0.43*	
Age	-0.23 0.02		0.00084*
Education	0.34 0.03		0.00075*
Sex	1.70 0.36		0.00010*
Lifetime alcohol use	-0.08 0.59		0.056
Current alcohol use	-0.08 0.59		0.056

Note : , parameter estimate ; SE(), standard error of parameter estimate ; R², variance explained by each model

Variables written in bold characters were included in the final model

* : $p<0.0001$

2(stage 2)
 가
 2
 (R²) 1

MMSE - KC

MMSE - KC

MMSE - KC 10.0%, 30.4%, 2.3%

4. 정상기준 자료산출 및 기준표 작성

MMSE - KC

(cell) 가
 60~74 , 65~79 , 70~84 , 75~90

(overlapping age table)

³⁵⁾
 (midpoint age) 67 , 72 , 77 , 82

Table 3. Mean, standard deviation, and selected percentiles of the MMSE-KC by age, educational level, and sex in the normal Korean elderly

Educational level(years)	Male			Female		
	0 - 3	4 - 6	7	0 - 3	4 - 6	7
Age (years)						
60 - 74*						
N	20	40	65	92	137	65
Mean	24.9	27.3	28.1	23.7	26.4	28.2
Sd	2.7	2.2	1.7	3.5	2.6	1.5
5th Percentile	18.0	23.1	24.3	16.0	20.9	25.0
Lower quartile	23.0	25.0	27.0	22.0	25.0	27.0
Median	25.0	27.5	29.0	24.0	27.0	29.0
Upper quartile	26.0	29.0	29.0	26.0	28.0	29.0
65 - 79**						
N	30	53	81	120	131	67
Mean	24.6	26.9	28.0	22.5	26.2	27.7
Sd	3.3	2.4	1.7	3.9	2.8	1.9
5th Percentile	17.5	23.0	24.1	15.0	20.5	24.2
Lower quartile	23.0	25.0	27.0	20.0	24.0	26.0
Median	25.0	27.0	28.0	23.0	27.0	28.0
Upper quartile	26.0	29.0	29.0	25.0	28.0	29.0
70 - 84***						
N	31	43	56	97	64	42
Mean	24.5	26.3	27.9	21.2	25.9	27.0
Sd	3.2	2.7	1.7	4.1	2.9	2.1
5th Percentile	17.0	21.2	24.0	14.9	20.0	23.0
Lower quartile	23.0	24.0	27.0	18.0	24.0	25.8
Median	25.0	26.0	28.0	22.0	26.0	27.0
Upper quartile	26.3	28.0	29.0	24.0	28.0	29.0
75 - 90****						
N	25	28	31	72	22	21
Mean	23.7	25.5	27.2	19.5	24.9	26.2
Sd	4.3	2.9	1.6	4.1	3.0	2.2
5th Percentile	13.5	20.5	24.0	13.7	20.0	20.0
Lower quartile	22.3	23.3	25.5	16.0	23.0	25.0
Median	24.5	25.0	27.0	19.0	24.0	26.0
Upper quartile	26.8	28.0	28.0	22.5	27.5	28.0

* : Normative data from this age range are used for persons whose ages range from 60 to 69 years

** : Normative data from this age range are used for persons whose ages range from 70 to 74 years

*** : Normative data from this age range are used for persons whose ages range from 75 to 79 years

**** : Normative data from this age range are used for persons whose ages range from 80 to 90 years

가 . 4
 60~74 , 65~79 , 70~84 , 75~90 , (0~3) ,
 60~69 , (4~6) , (7) .
 70~74 , 75~79 , 80~90 3 MM-

, Battersby⁴⁵⁾ (retention) 가 가

, Kuusisto¹⁸⁾ ,⁵⁶⁾ .

19-21)46-51) MMSE - KC 가

가 DSM - (protective effect)가²³⁾²⁴⁾⁵⁷⁾

가⁵⁸⁾⁵⁹⁾

가⁶⁰⁾

가 MMSE - KC

MMSE, Vanhanen⁵²⁾

가 가 MMSE - KC 가 Pauer³⁵⁾

가 60~74 75~90

가 가 74 75 1~2¹⁶⁾

가 가 가 가

⁵³⁾⁵⁴⁾ 가 Mayo's

Older Americans Normative Studies(MOANOS project) , MM-SE - KC

61 - 65)

가
가 MM-SE - KC 5

ADRDA NINCDS - 5.9% 5%

()
(percentile) 가

34) 5

MMSE
60~69 0~3 , 4~6 , 7
17/18 , 22/23 , 23/24 ,
15/16, 20/21, 24/25가

MMSE 가
23/24 , 1)

가
(false positive rate) 65 5% 80
가 2 20% , 8)

1
2 가
가 가

가

5 MMSE NCDS - ADRDA NI-
(normal control group) , 34)
' Aging -
Associated Cognitive Decline(AACD) '

MMSE - KC
(relatively healthy sample) , 67)

가 MMSE - KC 가
가

kappa , kappa
0.63 . Landis & Koch⁶⁶⁾가 가
kappa 0.61~0.80 , 48.9% 60
(Substantial)

- in the elderly. *J Am Geriatr Soc* 33 : 228-235
- 4) Crum RM, Anthony JC, Bassett SS, Folstein MF (1993) : Population-based norms for the Mini-Mental State Examination by age and educational level. *JAMA* 269 : 2386-2391
 - 5) 권용철, 박종한(1989) : 노인용 한국판 Mini-mental state examination(MMSE-K)의 표준화 연구. *신경정신의학* 28 : 125-135
 - 6) 강연욱, 나덕렬, 한승혜(1997) : 치매환자들을 대상으로 한 K-MMSE의 타당도 연구. *대한신경과학회지* 15 : 300-307
 - 7) Park JH, Ko HJ, Park YN, Jung CH(1994) : Dementia among the elderly in a rural Korean community. *Br J Psychiatr* 164 : 796-801
 - 8) Woo JI, Lee JH, Yoo KY, Kim CY, Kim YI, Shin YS (1998) : Prevalence estimation of dementia in a rural area of Korea. *J Am Geriatr Soc* 46 : 983-987
 - 9) Anthony JC, LeResche L, Niaz U, Von Korff MR, Folstein MF(1982) : Limits of the 'Mini-Mental State' as a screening test for dementia and delirium among hospital patients. *Psychol Med* 12 : 397-408
 - 10) Escobar JI, Burnam A, Karno M, Forsythe A, Landsverk J, Golding JM(1986) : Use of Mini-Mental State Examination(MMSE) in a community population of mixed ethnicity : Cultural and linguistic artifact. *J Nerv Ment Dis* 174 : 607-615
 - 11) Fillenbaum GG, Hughes DC, Heyman A, George LK, Blazer DG(1988) : Relation of health and demographic characteristics to Mini-Mental State Examination score among community residents. *Psychol Med* 18 : 719-726
 - 12) Kittner SJ, White LR, Farmer ME, Wolz M, Kaplan E, Moes E, Brody JA, Feinleib M(1986) : The problem of education adjustment. *J Chron Dis* 39 : 163-170
 - 13) O'Connor DW, Pollitt PA, Treasure FP, Brook CPB, Reiss BB, Roth M(1989) : The influence of education, social class and sex on Mini-Mental State scores. *Psychol Med* 19 : 771-776
 - 14) 우종인, 이정희, 홍진표(1996) : 농촌지역 거주노인의 연령, 성별, 교육수준이 한국판 MMSE 점수에 미치는 영향에 관한 연구. *신경정신의학* 35 : 122-132
 - 15) 여형용, 이강욱, 이근희, 주진형, 김기웅, 이정희, 하규섭, 정도연, 김용식, 우종인(1998) : 도시 및 농촌 노인의 한국판 MMSE 점수에 사회인구학적 변인이 미치는 영향에 관한 연구. *신경병리학* 7 : 72-80
 - 16) Lezak MD(1995) : Neuropsychological assessment. New York, Oxford University Press. 288-312
 - 17) Elias MF, Wolf PA, D'Agostino RB, Cobb J, White LR(1993) : Untreated blood pressure is inversely related to cognitive functioning : the Framingham study. *Am J Epidemiol* 138 : 353-64
 - 18) Kuusisto J, Koivisto k, Mykkanen L, Helkala EL, Vanhanen M, Hanninen T, Pyorala K, Riekkinen P, Laakso M(1993) : Essential hypertension and cognitive function. The role of hyperinsulinemia. *Hypertension* 22 : 771-9
 - 19) Scott RD, Kritz-Silverstein,D, Barrett-Connor E, Wiederhold WC(1998) : The association of non-insulin-dependent diabetes mellitus and cognitive function in an older cohort. *J Am Geriatr Soc* 46 : 1217-1222
 - 20) U'ren RC, Riddle MC, Lezak MD, Bennington-Daivs M(1990) : The mental efficiency of the elderly person with type II diabetes mellitus. *J Am Geriatr Soc* 38 : 505-510
 - 21) Perlmutter LC, Hakami MK, Hodgson-Harrington C, Ginsberg J, Katz J, Singer DE, Nathan DM(1984) : Decreased cognitive function in aging non-insulin-dependent diabetic patients. *Am J Med* 77 : 1043-1048
 - 22) Graves AB, van Duijn CM, Chandra V, Graves AB, van Duijn CM, Chandra V, Fratiglioni L, Heyman A, Jorm AF, Kokmen E, Kondo K, Mortimer JA, Rocca WA (1991) : Alcohol and tobacco consumption as risk factors for Alzheimer disease : a collaborative reanalysis of case control studies. EURODEM Risk Factor Research Group. *Int J Epidemiol* 20 : S48-S57
 - 23) Lee PN(1994) : Smoking and Alzheimer's disease : a review of the epidemiological evidence. *Neuroepidemiol* 13 : 131-144
 - 24) Feinstein AR(1974) : Clinical biostatistics, XXV : a survey of the statistical procedures in general medical journals. *Clin Pharmacol Ther* 15 : 97-107
 - 25) Heeren TJ, Lagaay AM, Beek VCAv, Rooymans HGM, Hijmans W(1990) : *J Am Geriatr Soc* 38 : 1093-1096
 - 26) Bravo G, Hebert R(1997) : Age- and education-specific reference values for the Mini-Mental and modified Mini-Mental State Examinations derived from a non-demented elderly population. *Int J Geriatr Psychiatr* 12 : 1008-1018
 - 27) Dufouil C, Clayton D, Brayne C, Chi LY, Dening TR, Paykel ES, O'Connor DW, Ahmed A, McGee MA, Huppert FA(2000) : Population norms for the MMSE in the very old. *Neurology* 55 : 1609-1613
 - 28) Flynn JR(1987) : Massive IQ gains in 14 nations : What IQ tests really measure. *Psychological Bulletin* 101 : 171-191

- 29) 통계청 (1999) : 인구통계연감. 서울.
- 30) Lee JH, Lee KU, Lee DY, Kim KW, Jho JH, Lee KH, Kim SY, Han SH, Woo JI (2002) : Development of the Korean version of the Consortium to Establish a Registry for Alzheimer's Disease (CERAD) Assessment Packet (CERAD-K) : Clinical and Neuropsychological Assessment Batteries. *J Gerontol Psychol Sci* 57(1) : 47-53
- 31) 이동영, 윤종철, 이강욱, 주진형, 김기웅, 이정희, 우종인 (1999) : 치매 선별을 위한 한국어판 간이 Blessed 검사의 신뢰도 및 타당도. *신경정신의학* 38 : 1297-1307
- 32) American Psychiatric Association (1994) : Diagnostic and Statistical Manual of Mental disorders, 4th edition. Washington DC.
- 33) Williams GH (1988) : Approach to the patient with hypertension. In Fauci AS, Braunwald E, Isselbacher KJ, Wilson JD, Martin JB, Kasper DL, Hauser SL, Longo DL Eds., *Harrison's Principles of Internal Medicine*, 14 edition. New York, McGraw-Hill, pp202-204
- 34) McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM (1984) : Clinical Diagnosis of Alzheimers Disease : report of the NINCDS-ADRDA Work Group under the auspices of the Department of Health and Human Service Task Force on Alzheimers Disease. *Neurology* 34 : 939-944
- 35) Pauker JD (1988) : Constructing overlapping cell tables to maximize the clinical usefulness of normative data : Rationale and an example from neuropsychology. *J Clin Psychol* 44 : 930-933
- 36) Scherr PA, Alert MS, Funkenstein HH, Cook NR, Hennekens CH, Branch LG, White LR, Taylor JO, Evans DA (1988) : Correlates of cognitive function in an elderly community population. *Am J Epidemiol* 128 : 1084-1101
- 37) Welsh KA, Butters N, Mohs RC, Beekly D, Edland S, Fillenbaum G, Heyman A (1994) : The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part V. A normative study of the neuropsychological battery. *Neurology* 44(4) : 609-614.
- 38) Ganguli M, Ratcliff G, Huff FJ, Belle S, Kancel MJ, Fischer L, Seaberg EC, Kuller LH (1991) : Effects of age, gender, and education on cognitive tests in a rural elderly community sample : Norms from the Monongahela Valley independent Elders Survey. *Neuroepidemiol* 10 : 42-52
- 39) Petersen RC, Smith G, Kokmen E, Ivnik RJ, Tangalos EG (1992) : Memory function in normal aging. *Neurology* 42 : 396-401
- 40) Blecker M, Bolla-Wilson K, Agnew J, Meyers D (1988) : Age-related sex differences in verbal memory. *J Clin Psychology* 44 : 403-411
- 41) Mungas D, Marshall S, Weldon M, Haan M, Reed B (1996) : Age and education correction of mini-mental state examination for English and Spanish-speaking elderly. *Neurology* 46 : 700-706
- 42) Magaziner J, Basset SS, Hebel JR (1987) : Predicting performance on the Mini-Mental State Examination : Use of age and education-specific equations. *J Am Geriatr Soc* 35 : 996-1000
- 43) Liu HC, Teng EL, Lin KN, Hsu TC, Guo NW, Chou P, Hu HH, Cheng WN, Chiang BN (1994) : Performance on a dementia screening test in relation to demographic variables. *Arch Neurol* 51 : 910-915
- 44) Farmer ME, Kittner SJ, Abbott RD, Wolz MM, Wolf PA, White LR (1990) : Longitudinally measured blood pressure, antihypertensive medication use, and cognitive performance : the Framingham study. *J Clin Epidemiol* 43 : 475-480
- 45) Battersby C, Hartley K, Fletcher AE, Markowe HJ, Brown RG, Styles W, Carne S, Jamieson T, Koppel I, Fraser S (1993) : Cognitive function in hypertension : a community based study. *J Hum Hypertens* 7(2) : 117-23
- 46) Gregg EW, Yaffe K, Cauley JA, Rolka DB, Blackwell TL, Narayan KMV, Cummings SR (2000) : Is diabetes associated with cognitive impairment and cognitive decline among older women? the Study of Osteoporotic Fractures Research Group. *Arch Intern Med* 160 : 174-180
- 47) Strachan MWJ, Deary IJ, Ewing FME, Friera BM (1997) : Is type II diabetes associated with an increased risk of cognitive dysfunction? *Diabetes Care* 20 : 438-445
- 48) Elias PK, Elias MF, D'Agostino RB, Cupples LA, Wilson PW, Silbershatz H, Wolf PA (1997) : NIDDM and blood pressure as risk factors for poor cognitive performance : the Framingham Study. *Diabetes Care*. 20 : 1388-1395
- 49) Ott A, Stolk RP, Hofman A, van Harskamp F, Grobbee DE, Breteler MMB (1996) : Association of diabetes mellitus and dementia : the Rotterdam Study. *Diabetologia*. 39 : 1392-1397
- 50) Lowe LP, Tranel D, Wallace RB, Welty TK (1994) : Type II diabetes and cognitive function : a population-based study of native Americans. *Diabetes Care* 17 : 891-896

- 51) Reaven GM, Thompson LW, Nahum D, Haskins E (1990) : relationship between hyperglycemia and cognitive function in older NIDDM patients. *Diabetes Care* 13 : 16-21
- 52) Vanhanen M, Kuusisto J, Koivisto K, Mykkinen Lk, Helkala E-L, Hänninen T, Riekkinen Sr P, Soininen H, Laakso M (1999) : Type-2 diabetes and cognitive function in a non-demented population. *Acta Neurol Scand* 100 : 97-101
- 53) Grant I (1987) : Alcohol and the brain : Neuropsychological correlates. *J Consult Clin Psychol* 55 : 310-324
- 54) Parsons OA (1986) : Cognitive functioning in sober social drinkers : A review and critique. *J Stud Alcohol* 47 : 101-114
- 55) Baum-Baicker C (1985) : The psychological benefits of moderate alcohol consumption : a review of the literature. *Drug Alcohol Depend* 15 : 305-322
- 56) Elwood PC, Gallacher JEJ, Hopkinson CA, Pickering J, Rabbitt P, Stollery B, Brayne C, Huppert FA, Bayer A (1999) : Smoking, drinking, and other life style factors and cognitive function in men in the Caerphilly cohort. *J Epidemiol Community Health* 53 : 9-14
- 57) Letenneur L, Dartigues JF, Commenges D, Barberger-Gateau P, Tessier JF, Orgogozo JM (1994) : Tobacco consumption and cognitive impairment in elderly people : a population-based study. *Ann Epidemiol* 4 : 449-454
- 58) Hebert LE, Scherr PA, Beckett LA, Funkenstein HH, Albert MS, Chown MJ, Evans DA (1992) : Relation of smoking and alcohol consumption to incident Alzheimer's disease. *Am J Epidemiol* 135 : 347-355
- 59) Ford AB, Mefrouche Z, Friedland RP, Debanne SM (1996) : Smoking and cognitive improvement : a population-based study. *J Am Geriatr Soc* 44 : 905-909
- 60) Riggs JE (1996) : the "protective" influence of cigarette smoking on Alzheimer's and Parkinson's disease : quagmire or opportunity for neuroepidemiology : *Neurol Clin* 14 : 353-358
- 61) Ivnik RJ, Malec JF, Smith GE, Tangalos EG, Petersen RC (1996) : Neuropsychological tests' norms above 55 : COWAT, BNT, MAE Token, WRAT-R reading, AMNART, STROOP, TMT, and JLO. *Clin Neuropsychologist* 10 : 262-278
- 62) Ivnik RJ, Malec JF, Smith GE, Tangalos EG, Petersen RC, Kokmen E, Kurland LT (1992) : Mayo's Older Americans Normative Studies : Updated AVLT norms for ages 56-97. *Clin Neuropsychologist* 6 (suppl) : 83-104
- 63) Ivnik RJ, Malec JF, Smith GE, Tangalos EG, Petersen RC, Kokmen E, Kurland LT (1992) : Mayo's Older Americans Normative Studies : WAIS-R norms for ages 56-97. *Clin Neuropsychologist* 6 (suppl) : 1-30
- 64) Ivnik RJ, Malec JF, Smith GE, Tangalos EG, Petersen RC, Kokmen E, Kurland LT (1992) : Mayo's Older Americans Normative Studies : WMS-R norms for ages 56-94. *Clin Neuropsychologist* 6 (suppl) : 49-82
- 65) Paolo AM, Tr Ster AI, Ryan JJ (1997) : California Verbal Learning Test : Normative data for the elderly. *J Clin Exp Neuropsychol* 2 : 220-234
- 66) Landis J, Koch GG (1977) : The measurement of observer agreement for categorical data. *Biometrics* 33 : 159-174
- 67) Levy R (1994) : Working party of the International Psychogeriatric Association in collaboration with the World Health Organization Aging-Associated Cognitive Decline. *Int Psychogeriatr* 6(1) : 63-68

A Normative Study of the Mini-Mental State Examination in the Korean Elderly

**Dong Young Lee, M.D., Ph.D., Kang Uk Lee, M.D., Ph.D., Jung Hie Lee, M.D., Ph.D.,
Ki Woong Kim, M.D., Ph.D., Jin Hyeong Jhoo, M.D., Jong Chul Youn, M.D.,
Sung Youn Kim, M.D., Ph.D., Sung Il Woo, M.D., Ph.D., Jong Inn Woo, M.D., Ph.D.**

Department of Neuropsychiatry, Kyunggi Provincial Hospital for the Elderly, Yongin

This study aimed to determine the variables significantly related to the MMSE in the Korean version of the CERAD assessment battery (MMSE-KC) and present the normative information of the test in the Korean elderly.

The MMSE-KC was administered in a standardized manner to 618 healthy volunteers aged 60 to 90 who were recruited from 4 dementia clinics, 2 elderly welfare centers and 1 community population. The elderly with serious neurological, medical and psychiatric disorders including dementia and major depression were excluded through clinical evaluation using CERAD-K clinical assessment protocol. Univariate and multivariate analyses were performed to determine the variables significantly related to the MMSE-KC score.

Age, educational level, and sex were found to have statistically significant effect on the test score. Based on this result, overlapping age normative table (60 to 74, 65 to 79, 70 to 84, and 75 to 90 years of age) with 3 educational strata (0 to 3 years, 4 to 6 years, and 7 years and more) separately for both male and female were developed. For resulting 24 normative units, 5 percentile, lower quartile, median, upper quartile, mean and standard deviation were calculated.

The normative data from this study can be widely used as reference values to objectively interpret the MMSE-KC scores of the Korean elderly.

KEY WORDS : MMSE-KC · MMSE · Norm · Age · Education · Sex.

□ 부 록 □

MMSE-KC
Mini-Mental Status Examination
in the Korean Version of the CERAD Assessment Packet

검사자는 “ 가 가 가 ”라는 말로 검사를 시작한다.

1.	?	0	1	9
2.	?	0	1	9
3.	?	0	1	9
4.	?	0	1	9
5.	? [피검자가 음력을 사용하면 음력으로 묻는다.]	0	1	9
6.	가 / / ?	0	1	9
7.	/ / ?	0	1	9
8.	/ / ?	0	1	9
9.	?	0	1	9
10.	?	0	1	9

11.	가 가 가 가 가			
	가			
	[문항 11은 첫 응답으로만 평가한다. 첫 응답에서 물건의 이름을 모두 말하지 못하는 경우는 문항 13의 지연 회상을 정확하게 검사하기 위해 '물건 이름을 불러주고 기억하도록 하는 과정'을 3회까지 반복할 수 있다.]			
	_____	0	1	9
	_____	0	1	9
	_____	0	1	9

12.	가			
	[피검자가 글자를 순서대로 바르게 말할 수 있도록 필요하다면 이 단어를 몇 차례 반복하여 말해 줄 수 있다.]			
	[피검자의 답을 위의 네모 칸에 기록한다. 점수는 뒷편에 제시한 기준에 따라 계산하여 우측 네모 칸에 기록한다.]			

